

SEMINÁRIO

LÓGICA MATEMÁTICA

2 de Dezembro | 16h00 | sala 6.2.33

A new cohomology for algebraic varieties over non-achimedean fields (*part II*)

Mário Edmundo
(FCUL e CMAF_cIO, Universidade de Lisboa)

Abstract:

The interplay between analytic geometry over non-archimedean fields and tropical geometry is a very active area with several applications in fields such as algebraic and arithmetic geometry. Recently, Hrushovski and Loeser introduced a model-theoretic account of the Berkovich's analytification of algebraic varieties: given a variety V over a non-archimedean field K , Hrushovski and Loeser associated to V the space \hat{V} , the stable completion of V , and showed a very deep connection between V and the tropical semi-group Γ_{∞} where Γ is the value group of K : there is a deformation retraction from \hat{V} to a definable subset of some finite power of Γ_{∞} . An analogous result was earlier proved by Berkovich for V^{an} under strong algebraic restrictions on the variety V . In this talk we report on the ongoing work (with P. Kovácsics and J. Ye) where we develop a sheaf cohomology theory for the spaces \hat{V} . When the field K is maximally complete of rank one, the spaces \hat{V} and $|V^{\text{an}}|$ are naturally homeomorphic and we recover results proved by Berkovich for the cohomology groups.