

SEMINÁRIO

ANÁLISE E EQUAÇÕES DIFERENCIAIS

12 de Dezembro | 14h30 | sala 6.2.38

A phase-field approximation for a surface-evolution energy with a density

Marco Caroccia
(Scuola Normale Superiore & Università di Firenze)

Abstract:

A perimeter type energy leading an epitaxial growth process is treated. We consider an energy of the type

$$\mathcal{F}(E, u) = \int_{\partial E} \psi(u)^{n-1}$$

where a density $u: \partial E \rightarrow \mathbb{R}$ living on the boundary of E is considered as a variable of the problem.

Such energy has been used to model an epitaxial growth process with the presence of adatoms: free atoms of the crystal allowed to move on the surface and which impacts the chemical potential of the evolution. In this context the set E represents the bulk, ∂E is the surface that is evolving in the process and the density u catches the presence of the so-called adatoms.

Such energy lacks of lower-semicontinuity in the natural topology under which the problem can be considered. In a contribution of 2016 jointly with Riccardo Cristoferi and Laurent Dietrich we provided a lower-semicontinuous envelope of \mathcal{F} and, subsequently, a phase-field approximation is developed.

I will briefly describe the models involving \mathcal{F} , the state of art and the lower-semicontinuous envelope calculation.

Financiado por Fundos Nacionais através da FCT – Fundação para a Ciência e a Tecnologia no âmbito do projeto UID/MAT/04561/2019

