

# SEMINÁRIO

## ANÁLISE E EQUAÇÕES DIFERENCIAIS

**26 de Julho | 13h30 | sala 6.2.33**

EXISTENCE AND DECAY RATES OF  $L^2$  NORMS FOR A GENERALIZED SEMILINEAR DISSIPATIVE EQUATION OF BOUSSINESQ/PLATE TYPE.

Ruy Coimbra Charão  
(Federal University of Santa Catarina)

ABSTRACT. we study existence, uniqueness and asymptotic behavior of solutions to the following Cauchy problem for a generalized second order semilinear equation of Boussinesq/Plate type under effects of a fractional dissipation

$$\begin{cases} u_{tt} + (-\Delta)^\delta u_{tt} + (-\Delta)^\alpha u + (-\Delta)^\theta u_t = \beta(-\Delta)^\gamma(f(u)), \\ u(0, x) = u_0(x), \\ u_t(0, x) = u_1(x), \end{cases} \quad (1)$$

where  $\beta \neq 0$  is a real constant,  $u = u(t, x)$  with  $(t, x) \in (0, \infty) \times \mathbb{R}^n$  and the exponents of the Laplacian operators are constants satisfying  $0 \leq \delta \leq \alpha$ ,  $0 \leq \theta \leq \frac{\alpha + \delta}{2}$  and  $\min\{0, \frac{\alpha}{2} - \frac{n}{4}\} \leq \gamma \leq \frac{\alpha + \delta}{2}$ . The nonlinearity  $f$  behaves as  $f(s) = s^p$  with  $p > 1$ . To show that the decay rates are optimal we use an asymptotic expansion of the solution in the Fourier space.